

WORLD INTELLECTUAL PROPERTY ORGANION International Bureau



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification 5:	A1	(11) International Publication Number: WO 91/18
C07D 401/12	AI	(43) International Publication Date: 12 December 1991 (12.12
(21) International Application Number: PCT/SI (22) International Filing Date: 5 June 1991 (30) Priority data: 9002043-9 7 June 1990 (07.06.90) (71) Applicant: AKTIEBOLAGET ASTRA [SE/SE]: Södertälje (SE). (72) Inventor: BRÄNDSTRÖM, Arne, Elof; Andersonsgatan 13 B, S-415 06 Göteborg (SE). (74) Agents: DANIELSSON, Sten et al.; AB Astra, F. partment, S-151 85 Södertälje (SE).	(05.06. ; S-151 ers Mat	(European patent), BF (OAPI patent), BG, BJ (O patent), BR, CA, CF (OAPI patent), CG (OAPI patent), CH, CH (European patent), CI (OAPI patent), (OAPI patent), DE, DE (European patent), DK, (European patent), ES, ES (European patent), FI, (European patent), GA (OAPI patent), GB, GB (E pean patent), GN (OAPI patent), GR (European tent), HU, IT (European patent), JP, KP, KR, LK, LU (European patent), MC, MG, ML (OAPI patent), MR (OAPI patent), MW, NL, NL (European patent), NO, PL, RO, SD, SE, SE (European patent), SN (O patent), SU, TD (OAPI patent), TG (OAPI patent).

(54) Title: IMPROVED METHOD FOR SYNTHESIS


(57) Abstract

The present invention relates to an improved method for the synthesis of omeprazole, comprising the steps of reacting 5-methoxy-2-[(4-methoxy-3,5-dimethyl-2-pyridinyl)-methylthio]-1H-benzimidazole with m-chloroperoxy-benzoic acid in a methylene chloride solution at a substantially constant pH of about 8.0 to 8.6; extracting the reaction mixture with aqueous NaOH; separating the aqueous phase from the organic phase; and adding an alkyl formate to the aqueous phase, resulting in crystallization of omeprazole.

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	ES	Spain _	MG	Madagascar
AU	Australia	FI	Finland	ML	Mali
BB	Barbados	FR	France	MN	Mongolia
BE	Belgium	GA	Gabon	MR	Mauritania
BF	Burkina Faso	GB	United Kingdom	MW	Malawi
BG	Bulgaria	GN	Guinca	NL	Netherlands
BJ	Benin	GR	Greece	NO	Norway
BR	Brazil	HU	Hungary	PL	Poland
CA	Canada	IT	Italy	RO	Romania
CF	Central African Republic	JP	Japan	SD	Sudan
CG	Congo	KP	Democratic People's Republic	SE	Sweden
CH	Switzerland		of Korea	SN	Senegal
CI	Côte d'Ivoire	KR	Republic of Korea	SU	Soviet Union
CM	Cameroon	LI	Liechtenstein	TD	Chad
CS	Czechoslovakia	LK	Sri Lanka	TG	Togo
DE	Germany	LU	Luxembourg	US	United States of America
DK	Denmark	MC	Monaco		





Improved method for synthesis

Technical field

5 The present invention relates to an improved method for the synthesis of 5-methoxy-2-[[(4-methoxy-3,5-dimethyl-2-pyridinyl)-methyl]sulfinyl-1H-benizimidazole, referred to under its generic name omeprazole throughout the following specification and claims.

10

Prior art

US-A-4 255 431 discloses a process for the synthesis of omeprazole comprising the steps of reacting 5-methoxy-2[(4-methoxy-3,5-dimethyl-2-pyridinyl)-methylthio]-1H-benzimidazole in a methylene chloride solution with m-chloroperoxybenzoic acid resulting in the formation of omeprazole and m-chlorobenzoic acid. omeprazole is highly sensitive to acids, and the reaction mixture has to be maintained at a low temperature to prevent excessive decomposition in the reaction mixture.


The product is worked-up by filtering-off of m-chloro-benzoic acid formed during the reaction. The filtrate is diluted with methylene chloride, is extracted with Na₂CO₃ solution, dried and evaporated. The resulting omeprazole product is contaminated with starting materials and by-products.

30 Summary of the invention

The object of the present invention is to provide an improved method for the synthesis of omeprazole, which eliminates the drawbacks of previously known methods.

35

This object is achieved according to the present inven-



tion, which is characterized by the steps of reacting 5methoxy-2-[(4-methoxy-3,5-dimethyl-2-pyridinyl)-methylthio]-1H-benzimidazole (below denoted Compound I) with mchloroperoxybenzoic acid in a methylene chloride solution
at a substantially constant pH of about 8.0 to 8.6;
extracting the reaction with aqueous NaOH; separating the
aqueous phase from the organic phase; and adding an alkyl
formate to the aqueous phase, resulting in crystallization
of omegrazole.

10

The m-chloroperoxybenzoic acid is suitably used in an amount of 0.7 - 1.4 molar equivalents of Compound I, and preferably in an amount of 0.9 - 1.2 molar equivalents.

15 According to one embodiment of the invention, the alkyl formate is methylformate or ethylformate, methylformate being preferred.

The alkyl formate is suitably used in an amount of 1.2 - 2.0 molar equivalents of Compound I, and preferably in an amount of 1.5 - 1.8 molar equivalents.

One important feature of the method according to the invention is that unreacted sulfide is not transferred into the aqueous phase upon the extraction with aqueous NaOH. Another important feature is that m-chlorobenzoic acid does not crystallize upon the addition of methylformate to the aqueous phase, thereby eliminating the need of filtering-off of m-chlorobenzoic acid in a previous step.

The pH of the reaction mixture may be maintained within the pH range of 8.0 - 8.6 with the aid of pH static titration with NaOH or with the use of a buffer. Preferred buffers are sodium bicarbonate and potassium bicarbonate.



A great advantage of the method according to the invention is that the reaction takes place in the organic methylene chloride phase while the m-chlorobenzoic acid formed during the reaction goes into the aqueous phase containing the buffer, in the case a buffer is used. Because of this, omeprazole formed does not stay in contact with the acid and the reaction may be performed at a temperature above 0°C.

10 According to one embodiment of the invention the pH of the aqueous NaOH phase is kept at above about 12.

According to another embodiment of the invention the crystallization of omeprazole is performed at a pH of above 9.

The invention will be further illustrated below with a non-limiting example.

20 Example

5-methoxy-2-[(4-methoxy-3,5-dimethyl-2-pyridinyl)methylthio]-1H-benzimidazole (16.2 g; 0.0492 mol) is
reacted with m-chloroperoxybenzoic acid (13.6 g; 0.0537

25 mol) in CH₂Cl₂ acting as a solvent at a pH of 8.6, which
is maintained by the presence of KHCO₃ (5.6 g; 0.056 mol)
acting as a buffer. The temperature is maintained at about
0°C during the addition.

30 Diluted NaOH is added to a pH above 12 and the ${\rm CH_2Cl_2}$ phase is separated off.

Methylformate (4.7 g) is charged to the water phase and the pH is kept above 9, whereupon omeprazole crystallizes.

35 The crystals are filtered off and are washed with water and methanol at a temperature of about 0°C. The washed crystals are dried under vacuum. Yield: 15.6 g (92 %).



35

Claims

- 1. An improved method for the synthesis of omeprazole, c h a r a c t e r i z e d by the steps of reacting 55 methoxy-2-[(4-methoxy-3,5-dimethyl-2-pyridinyl)-methyl-thio]-1H-benzimidazole (Compound I) with m-chloroperoxy-benzoic acid in a methylene chloride solution at a substantially constant pH of about 8.0 to 8.6; extracting the reaction mixture with aqueous NaOH; separating the aqueous phase from the organic phase; and adding an alkyl formate to the aqueous phase, resulting in the crystallization of omeprazole.
- 2. Method according to claim 1, c h a r a c t e r i -15 z e d in that the m-chloroperoxybenzoic acid is used in an amount of 0.7 - 1.4, preferably 0.9 - 1.2, molar equivalents of Compound I.
- 3. Method according to claim 1 or 2, characte-20 rized in that the alkyl formate is methylformate.
- 4. Method according to claims 1 3, c h a r a c t e r i z e d in that pH of the reaction mixture is maintained within the pH range of 8.0 8.6 with the aid of pH 25 static titration with NaOH.
- 5. Method according to claims 1 4, characterized in that pH of the reaction mixture is maintained within the pH range of 8.0 8.6 with the use of a buffer.
 - 6. Method according to claim 5, characterized in that the buffer is sodium bicarbonate or potassium bicarbonate.
 - 7. Method according to claims 1 6, characterized in that the pH of the aqueous NaOH phase is



kept at above about 12.

- 8. Method according to claims 1 7, c h a r a c t e r i z e d in that the alkyl formate is added in an amount
 5 of 1.2 2.0, preferably 1.5 1.8, molar equivalents of Compound I.
- Method according to claims 1 8, c h a r a c t e r i z e d in that the crystallization of omeprazole is
 performed at a pH of above 9.





INTERNATIONAL SEARCH REPORT

			3E 91/00402
I. CLASSIFICA	TION OF SUBJECT MATTER (if several classi	fication symbols apply, indicate all) b	
According to Inte IPC5: C 07	ernational Patent Classification (IPC) or to both I D 401/12	National Classification and IPC	
II. FIELDS SEAF	RCHED		
	Minimum Docume	entation Searched 7	
Classification Syst	em	Classification Symbols	<u></u> .
100C	C 07 D		
IPC5	C 07 D		··-
		r than Minimum Documentation ts are Included in Fields Searched ⁸	
SE,DK,FI,NO	classes as above		•
III. DOCUMENTS	CONSIDERED TO BE RELEVANTS		
Category = C	itation of Document, ¹¹ with indication, where ap	propriate, of the relevant passages ¹²	Relevant to Claim No.13
	A, 4255431 (AKTIEBOLAGET HÄ 10 March 1981, see example 1	SSLE)	1
	see example 1		
	A, 4182766 (HOFFMANN-LA-ROC 8 January 1980, see example 17	HE INC)	1
		XCCI E)	. 1
	A1, 8705021 (AKTIEBOLAGET H 27 August 1987, see example 1	ASSLE)	
	A, 0197013 (AKTIEBOLAGET HÄ 8 October 1986, see example 2 	SSLE)	1
"A" document d	ories of cited documents; ¹⁰ efining the general state of the art which is not to be of particular relevance	"T" later document published after or priority date and not in conflicited to understand the principle invention	he international filing date of with the application but or theory underlying the
filing date	ment but published on or after the international	cannot be considered novel or c	e, the claimed invention annot be considered to
"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified) "Y" document of particular relection of the considered to in decument is combined with			or more other such docu-
other mean:		in the art.	onalogs to a belson served
	ublished prior to the international filing date bu he priority date claimed	"&" document member of the same	patent ramily
Date of the Actual (ON Completion of the International Search	Date of Mailing of this International Se	earch Report
28th August		1991 -09- 1 0	
International Searc	hing Authority	Signature of Authorized Officer	
	EDISH PATENT OFFICE econd sheet) (January 1985)	Göran Karlsson	

BEST AVAILABLE COPY





International Application No. PCT/SE 91/00402

ategory "	MENTS CONSIDERED TO BE RELEVANT (CONTINUED FROM THE SECOND SHEET) Citation of Document, with indication, where appropriate, of the relevant passages	Relevant to Claim No
	EP, A, 0163842 (F. HOFFMANN-LA ROCHE & CO.) 11 December 1985, see example 4	1
		· <u>-</u> ·
İ		
.		
ĺ		
-		
	·	
İ	·	
		•
		_
		•
	·	
1		
	.	
İ		
1	1	







ANNEX TO THE INTERNATIONAL SEARCH REPORT ON INTERNATIONAL PATENT APPLICATION NO.PCT/SE 91/00402

This annex lists the patent family members relating to the patent documents cited in the above-mentioned international search report. The members are as contained in the Swedish Patent Office EDP file on 91-06-27 The Swedish Patent Office is in no way liable for these particulars which are merely given for the purpose of information.

c	Patent document ited in search report	Publication date		family hber(s)	. Publication date
US-A-		81-03-10	AT-B- AT-B- AT-B- AT-B- AT-B- AU-B- AU-D- CA-A- CA-A- EP-A-B- JP-C- JP-C- JP-A- JP-B- JP-B- SE-A- US-A-	374471 374472 374473 375365 389995 529654 4602779 1127158 1129417 0005129 1312930 1504537 54141783 58192880 60034956 63053191 7804231 4337257 4508905	84-04-25 84-04-25 84-04-25 84-07-25 90-02-26 83-06-16 79-10-18 82-07-06 82-08-10 79-10-31 86-04-28 89-07-13 79-11-05 83-11-10 85-08-12 88-10-21 79-10-15 82-06-29 85-04-02
US-A-	4182766	80-01-08	AT-B- AU-B- AU-D- CA-A- DE-A- EP-A-B- FR-A-B- GB-A-B- JP-A- NL-A- OA-A- SE-A- US-A-	368152 519711 3981778 1106849 2840591 0001279 2403340 2004281 54059274 7809529 6054 7809792 4248880	82-09-27 81-12-17 80-03-20 81-08-11 79-03-29 79-04-04 79-04-13 79-03-28 79-05-12 79-03-21 81-06-30 79-03-20 81-02-03
WO-A1-	8705021	87-08-27	EP-A- JP-T-	0242341 63502658	87-10-21 88-10-06
EP-A-	0197013	86-10-08	JP-A-	61205211	86-09-11
EP-A-	0163842	85 -12- 11	AU-B- AU-D- AU-D- JP-A- SU-A-	589555 4122885 4136685 60233070 1362402	89-10-19 85-10-24 85-10-24 85-11-19 87-12-23







ANNEX TO THE INTERNATIONAL SEARCH REPORT ON INTERNATIONAL PATENT APPLICATION NO.PCT/SE 91/00402

This annex lists the patent family members relating to the patent documents cited in the above-mentioned international search report. The members are as contained in the Swedish Patent Office EDP file on 91-06-27 The Swedish Patent Office is in no way liable for these particulars which are merely given for the purpose of information.

c	Patent document ited in search report	Publication date		t family nber(s)	· Publication date
EP-A-	0163842	85-12-11	SU-A- US-A- US-A-	1396965 4634710 4981861	88-05-15 87-01-06 91-01-01